Acc. No.: TH-30

## **SUMMARY**

In summary, the present thesis, demonstrates that M150 costimulates CD8<sup>+</sup> T cells to mature into an effector cytotoxic T cells. The findings can briefly be summarized as follows:

- 1. Antibody raised against M150, in hamster, recognizes M150 on the surface of macrophages and LPS activated B cells but not on resting B cells.
- 2. This antibody inhibits CD8+ T cell response in MLR. Inhibition of MLR by anti-M150 varies and is determined by type of stimulator cells used. Blocking is maximum when macrophages are used as stimulator cells as compared to B cells. This signifies that M150 is a major costimulatory molecule as far as macrophages are concerned.
- 3. Isolated and liposomised M150 stimulates CD8+ T cells only in association with anti-CD3. This proves that it is a costimulatory molecule.
- 4. The activation of CD8+ T cells by M150 is mediated by signal transduction. Signaling through the TcR by anti-CD3 antibody or anti-TcR antibody in combination with M150 has a synergistic effect on the levels of calcium.
- 5. In response to M150, as costimulatory signal, a protein of 22-23 kDa is tyrosine phosphorylated. Costimulation also results in the down regulation of CD45RB expression.
- 6. CD8+ T cells acquire granazyme A and perforin as a result of costimulation by M150.

- 7. The CD8+ T cells maintain the expression of Fas L only in case they are stimulated with both first (anti-CD3) and second signal (M150).
- 8. The anti-CD3 activated CD8+ T cells express Fast valued with both first and second signal do not.
- 9. Anti-Fas antibodies fail to induce apoptosis is such T cells simultaneously express high

٦e