



Data in Brief

Genome sequencing and annotation of *Acinetobacter haemolyticus* strain MTCC 9819^T☆Indu Khatri^{b,1}, Nitin Kumar Singh^{a,1}, Srikrishna Subramanian^{b,*}, Shanmugam Mayilraj^{a,*}^a Microbial Type Culture Collection and Gene bank (MTCC), CSIR-Institute of Microbial Technology, Chandigarh 160036, India^b Protein Science and Engineering, CSIR-Institute of Microbial Technology, Chandigarh 160036, India

ARTICLE INFO

Article history:

Received 17 September 2013

Received in revised form 23 October 2013

Accepted 23 October 2013

Available online 27 November 2013

Keywords:

Acinetobacter haemolyticus strain MTCC 9819^T

Whole genome

Illumina-HiSeq 1000 technology

CLCbio wb6

Rapid annotations using subsystems technology (RAST)

ABSTRACT

The genus *Acinetobacter* consists of 31 validly published species ubiquitously distributed in nature and primarily associated with nosocomial infection. We report the 3.4 Mb genome of *Acinetobacter haemolyticus* strain MTCC 9819^T. The genome has a G + C content of 40.0% and includes 3 rRNA genes (5S, 23S, 16S) and 65 aminoacyl-tRNA synthetase genes.

© 2013 The Authors. Published by Elsevier Inc. All rights reserved.

Specifications

Organism/cell line/tissue	<i>Acinetobacter haemolyticus</i>
Strain(s)	MTCC 9819 ^T
Sequencer or array type	Sequencer; the Illumina-HiSeq 1000
Data format	Processed
Experimental factors	Microbial strain
Experimental features	Whole genome sequencing of <i>A. haemolyticus</i> strain MTCC 9819 ^T , assembly and annotation
Consent	n/a

Direct link to the data

Direct link: <http://www.ncbi.nlm.nih.gov/nucore/ASYX00000000>.

Genus *Acinetobacter* was proposed by Brisou and Pre'vot in 1954 [1]. This genus comprises Gram-negative, strictly aerobic, nonfermenting, nonfastidious, nonmotile, catalase-positive, oxidase-negative bacteria with a DNA G + C content of 39% to 47% [2]. According to Euzéby's list of prokaryotic names with standing in nomenclature (<http://www.bacterio.cict.fr/a/acinetobacter.html>) the genus *Acinetobacter* consists of 31 validly published species. *Acinetobacter haemolyticus* was proposed by Bouvet and Grimont 1986 [3]; it was isolated from human clinical specimens and environment, with characteristics corresponding to those of the genus *Acinetobacter*. The organism in this study is *Acinetobacter haemolyticus* strain MTCC 9819^T equivalent to DSM 6962 (= ATCC 17906, CIP 64.3, NCTC 10305).

A. haemolyticus strain MTCC 9819^T was obtained from MTCC and grown on tryptic soya agar medium (TSA; HiMedia) at 30 °C. Genomic DNA was extracted from 36 hour old culture using ZR Fungal/Bacterial DNA MiniPrep™ as per manufacturer's instructions. Identification and sequencing of 16S rRNA was performed as described by Mayilraj et al. 2006 [4]. To determine the phylogenetic relationship of strain MTCC 9819^T, the 16S rRNA sequence consisting of 1502 bp was compared with those of type strains of species of related genera and identification of phylogenetic neighbors and the calculation of pairwise 16S rRNA gene sequence similarities were achieved using the EzTaxon server [5] and aligned using mega version 5.0 [6]. Phylogenetic trees were constructed using the neighbor-joining algorithm. Bootstrap analysis was performed to assess the confidence limits of the branching (Fig. 1.).

The genome of *A. haemolyticus* strain MTCC 9819^T was sequenced using Illumina-HiSeq 1000 technology. Sequencing resulted in 5,755,416 paired-end reads (insert size of 350 bp) of length 101 bp. A total of 5,580,067 high-quality reads with approximately 166× coverage were assembled with CLCbio wb6 (word size = 50 and bubble size = 100) and to obtain 182 contigs (N₅₀, 57,666 bp) of 3,384,229 bp with an

☆ This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

* Corresponding authors at: Institute of Microbial Technology (IMTECH), Sector 39-A, Chandigarh 160036, India. Tel.: +91 172 6665166; fax: +91 172 2695215.

E-mail address: mayil@imtech.res.in (S. Mayilraj).

¹ Both are first authors, equally contributed.




Fig. 1. Phylogenetic tree constructed using the neighbor-joining algorithm, shows the position of *A. haemolyticus* strain MTCC 9819^T relative to the type strains of the other species within the genus *Acinetobacter*.

average GC content of 40%. The functional annotation was carried out by RAST (rapid annotation using subsystem technology) [7], Fig. 2 shows the subsystem distribution of strain *A. haemolyticus* strain MTCC 9819^T, and tRNA was predicted by tRNAscan-SE 1.23 [8] and rRNA genes by RNAmmer 1.2 [9]. The genome contains 3 rRNA genes (5S, 23S, 16S) and 65 aminoacyl-tRNA synthetase genes. A total of 3122 coding regions (1546 transcribed from the positive strand and 1576 from the negative

strand) were found in the genome, of which 2298 (74%) could be functionally annotated. The genome coding density is 87% with an average gene length of 911 bp. The annotated genome has 65 genes responsible for resistance to antibiotic and toxic compounds including 13 genes for MDR efflux pumps. One hundred and six genes code for membrane transport proteins. Forty four genes are involved in response to oxidative stress, 10 for osmotic stress response and 16 genes for heat




Fig. 2. Sub-system distribution of strain *A. haemolyticus* strain MTCC 9819^T (based on RAST annotation server).

shock and many more stress responses, all summed up to 94 genes for stress response are present.

The functional comparison of genome sequences available on the RAST server revealed the closest neighbors of *A. haemolyticus* strain MTCC 9819^T as *Acinetobacter junii* SH205 (score 503) followed by *Acinetobacter baumannii* ACICU (score 489), *Acinetobacter haemolyticus* ATCC 19194 (score 476) and *Acinetobacter baumannii* AB0057 (score 453).

Nucleotide sequence accession number

The *A. haemolyticus* strain MTCC 9819^T whole genome shot gun (WGS) project has been deposited at DDBJ/EMBL/GenBank under the project accession ASYX01000000 of project (01) which has the accession numbers ASYX01000000 and consists of sequences ASYX01000001 ASYX01000182.

Conflict of interest

The authors declare that there is no conflict of interest on any work published in this paper.

Acknowledgments

This work was funded by CSIR-IMTECH. N.K.S. is supported by a University Grant Commission (UGC) fellowship, and I.K. is supported by a research fellowship from CSIR. We thank the Sandor next-

generation genomics facility for the help in obtaining the genome sequence. This is IMTECH communication number 0107/2013.

References

- [1] J. Brisou, A.R. Prevot, Etudes de systematique bacterienne. X. Revision des especes reunies dans le genre *Achromobacter*. Ann. Inst. Pasteur 86 (1954) 722–728.
- [2] A.Y. Peleg, H. Seifert, D.L. Paterson, *Acinetobacter baumannii*: emergence of a successful pathogen. Clin. Microbiol. Rev. 21 (2008) 538–582.
- [3] P.J.M. Bouvet, P.A.D. Grimont, Taxonomy of the Genus *Acinetobacter* with the recognition of *Acinetobacter baumannii* sp. nov., *Acinetobacter haemolyticus* sp. nov., *Acinetobacter johnsonii* sp. nov., and *Acinetobacter junii* sp. nov. and emended descriptions of *Acinetobacter calcoaceticus* and *Acinetobacter lwoffii*. Int. J. Syst. Evol. Microbiol. 36 (2) (1986) 228–240.
- [4] S. Mayilraj, P. Saha, S. Korpole, H.S. Saini, *Ornithinimicrobium kibberense* sp. nov. isolated from the Himalayas, India. Int. J. Syst. Evol. Microbiol. 56 (2006) 1657–1661.
- [5] O. Kim, Y.J. Cho, K. Lee, S.H. Yoon, M. Kim, H. Na, S.C. Park, Y.S. Jeon, J.H. Lee, H. Yi, S. Won, J. Chun, Introducing EzTaxon-e: a prokaryotic 16S rRNA Gene sequence database with phylotypes that represent uncultured species. Int. J. Syst. Evol. Microbiol. 62 (2012) 716–721.
- [6] K. Tamura, D. Peterson, N. Peterson, G. Stecher, M. Nei, S. Kumar, MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Mol. Biol. Evol. 28 (2011) 2731–2739.
- [7] R.K. Aziz, D. Bartels, A.A. Best, M. DeJongh, T. Disz, R.A. Edwards, K. Formsma, S. Gerdes, E.M. Glass, M. Kubal, F. Meyer, G.J. Olsen, R. Olson, A.L. Osterman, R.A. Overbeek, L.K. McNeil, D. Paarmann, T. Paczian, B. Parrello, G.D. Pusch, C. Reich, R. Stevens, O. Vassieva, V. Vonstein, A. Wilke, O. Zagnitko, The RAST server: rapid annotations using subsystems technology. BMC Genomics 9 (2008) 75.
- [8] T.M. Lowe, S.R. Eddy, tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. Nucleic Acids Res. 25 (1997) 955–964.
- [9] K. Lagesen, P. Hallin, E.A. Rodland, H.H. Staerfeldt, T. Rognes, D.W. Ussery, RNAmmer: consistent annotation of rRNA genes in genomic sequences. Nucleic Acids Res. 35 (2007) 3100–3108.